



# Neutrinos: Physics Beyond the Standard Model Through Multi Messenger Astronomy and Lessons from IceCube

Avni V. Iyer

Leland High School, 6677, San Jose, CA, 95120, United States; avni.v.iyer@gmail.com

## Abstract

Neutrinos, a class of nearly massless elementary particles that interact with only the weak nuclear and gravitational forces, possess a remarkable ability to traverse enormous distances without interference. Originating from processes like nuclear fusion in stars, supernova explosions, supermassive blackholes and radioactive decay, neutrinos are unique in their capacity to journey through space largely undisturbed. This characteristic helps answer some of the most fundamental questions about the universe such as the inner workings of astrophysical bodies and potentially the matter-antimatter asymmetry in the universe. By reviewing data from IceCube, the world's largest optical Cherenkov telescope, this paper presents a critical evaluation of the information from a major identified source of astrophysical neutrinos: blazars. This paper also evaluates and discusses the importance of multi messenger astronomy and the need to go beyond the standard model to understand the complex nature of neutrino emissions, antineutrino characteristics, and blazars.

## Keywords

Physics and Astronomy; Atomic, Molecular and Optical Physics; Nuclear and Particle Physics; IceCube; Multi Messenger Astrophysics; Blazar; Neutrinos; TXS 0506+056

## Introduction

Neutrinos are part of the standard model, a theoretical framework for particle physics (Figure 1). The model consists of fermions (leptons and quarks), bosons, and the Higgs boson.<sup>1</sup> Fermions are matter-carrying particles made up of two subclasses: quarks and leptons.<sup>1</sup> The particles in the class of leptons include the electron, muon, and tau particles as well as the electron neutrino, muon neutrino, and tau neutrino. <sup>1</sup> Bosons on the other hand are force-carrying particles and include the photon, gluon, W boson, and Z boson.<sup>1</sup> The Higgs boson is a particle that creates the Higgs field, which fermions such as electrons and quarks interact with to obtain mass.<sup>1</sup>




Figure 1: A simplified representation showing matter carrying particles (leptons: yellow; quarks: green) and force-carrying particles (bosons: blue; Higgs boson: innermost circle). (Author)

## Neutrinos:

For neutrinos, the standard model has generated limited theories. Initially, scientists postulated, based on the standard model, that neutrinos possess no mass and consequently do not engage with the Higgs Boson in the same manner as other fermions. A second hypothesis was that neutrinos travel at the speed of light, in turn indicating that they are massless. However, Super-Kamiokande and the Sudbury Neutrino Observatory conducted experiments that prove neutrinos do have mass, though it is very close to zero, and were also shown to oscillate between 3 known flavors, the muon neutrino, tau neutrino, and electron neutrino respectively oscillating into a muon, tau, or electron upon interaction with matter.<sup>1</sup> These discoveries provide the foundation for our exploration of neutrinos from astrophysical sources.

Neutrinos are unique as they have a neutral charge, almost no mass, and travel close to the speed of light. Though neutrinos rarely interact with matter and only interact via the weak or gravitational forces, they are the most abundant particle with mass in the universe. They are produced during nuclear fission and fusion as well as during radioactive decay.<sup>2</sup>

Despite being the most abundant of all particles in the universe, detecting a neutrino is extremely rare. This makes them a challenge to study and for years the astrophysical sources of these high-speed neutrinos were not determined. Astrophysical neutrinos are of particular relevance due to their exceedingly high energies. These extreme energy neutrinos are produced only from interactions near supermassive black holes, which is why they provide insight to the unique processes within them.<sup>3</sup>

## CP Violation:

Another major question within the realm of neutrinos and astrophysics is the question of why the current universe has an abundance of matter over antimatter. Scientists' current model of the Big Bang includes matter and antimatter being formed in equal amounts. This also suggests that all the matter and antimatter annihilate each other, leaving nothing but light and radiation.

However, we know the universe contains an abundance of matter in comparison to antimatter. Charge-parity symmetry is when the charge can be inverted and coordinate directions mirrored without altering any other properties. Charge-parity violation, or CP violation, however, is a phenomenon where a particle and an antiparticle differ in their fundamental characteristics.<sup>4</sup> This is a key pathway to explain why matter can outweigh antimatter.

While scientists are exploring CP violation within quarks and their antiparticles, understanding neutrinos is also essential to this mystery. Their very low mass indicates that they might be the relatives of very heavy particles that formed near the Big Bang: undiscovered, right-handed particles that obtained mass from a method other than the Higgs field. Right-handed particles are those whose spin, or amount of angular momentum, aligns with their direction of motion. The disintegration of these particles may have led to CP violation that should be observed in neutrinos and antineutrinos today. The hypothesized process that neutrinos went through to cause an excess of matter over antimatter is known as leptogenesis.

By studying neutrinos and antineutrinos, scientists hope to understand more about CP violation and the matter-antimatter imbalance in the universe.

## Why Neutrinos are Key:

Particle physics encompasses 36 different fundamental particles, but neutrinos, and especially astrophysical neutrinos, are unique to our understanding of the universe.

Astrophysical neutrinos are released in the decay process of particles ejected by blazars. They are key in providing information about blazars because they only interact with gravity and the weak force, therefore traveling large distances without being affected by magnetic fields.<sup>5</sup> Astrophysical neutrinos provide a major source of information to answer key questions about the universe because of the large distance in space and time they travel. Once traced back to their sources, they give scientists information about processes that occur within blazars and characteristics of these cosmic bodies.

## Background

### Sources of Astrophysical Neutrinos:

Possible sources of cosmic neutrinos include black holes, pulsars, the remnants of supernovae, energetic explosions called gamma-ray bursts, and active galactic nuclei (AGNs).<sup>6</sup> Blazars are the main source of interest for this paper as they have been shown to be a primary source of astrophysical neutrinos.

Blazars are the active cores, or black holes, of giant galaxies. Because they are so bright, we can see them at very great distances. Thus, we can also see far back in time because of the amount of time it takes light to travel from a far away source to our detectors. The difference between blazars and other active galactic nuclei is that the jets of particles (cosmic rays) ricocheting from the blazar are directed towards Earth. These jets are formed as gasses around the active black hole rub against each other, losing kinetic energy in the process. This causes them to fall into the black hole and release visible heat and light. As the plasma moves faster, there is less space available, resulting in a crowding effect that

causes the plasma to ricochet away from the blackhole. Due to the irradiation of radio, optical, x-ray, gamma-ray, and neutrino emissions, blazars open the door for multiwavelength astrophysics.<sup>7</sup>

Two major subclasses of blazars have been identified: Flat-spectrum radio quasars (FSRQs) and Broad Line Lacertae Objects (BL Lac Objects). FSRQs are characterized by powerful GeV gamma rays and strong emission lines (lines of radiation produced by highly excited electrons). Emission lines are typically produced in the broad-line region (BLR) of a blazar, which is a region of dense gas and dust that is located close to the supermassive black hole at the center of the blazar. On the other hand, BL Lac Objects are less powerful with weaker emission lines, but the presence of Radiatively Inefficient Accretion Flows (RIAFs) near their nuclei cause them to be a point of intrigue for astrophysical neutrino sources, as is discussed later in the paper.


## Neutrino Fluxes - Photohadronic and Hadronuclear Interactions:

Understanding photohadronic and hadronuclear interactions that produce neutrinos is crucial to understanding more about the role of blazars in neutrino production covered later in this paper.


When cosmic rays from blazars interact with gasses or photons, they generate high-energy neutrino and gamma rays. Both hadronuclear (interaction with ambient matter in gasses) and photohadronic interactions (interaction with photons) produce charged and neutral pions (particles made up of a quark and antiquark) which play a significant role in the strong nuclear force.

Neutral pions decay into two gamma ray photons, contributing to the gamma-ray flux. Charged pions ( $\pi$ ) produce neutrinos ( $\nu$ ) along their decay chain. Neutrino flux refers to the rate at which neutrinos pass through a unit area. It's essentially a measure of the number of neutrinos per unit area, per unit time. Gamma rays are produced alongside high-energy neutrinos ( $\nu$  or  $\bar{\nu}$ ) and cascade down to lower energies. These gamma rays then lose energy over time, undergoing a process known as 'cascading,' ultimately resulting in the production of lower-energy photons.<sup>8</sup> These are explained by these equations:

Photo Hadronic interactions:




A high-energy proton ( $p$ ) collides with a photon ( $\gamma$ ), typically a gamma-ray photon. This interaction results in the creation of a neutron ( $n$ ) and a charged pion ( $\pi^+$ , if the original proton was positive, or  $\pi^-$ , if the proton was negative).




A charged pion ( $\pi^+$ ) or ( $\pi^-$ ) undergoes a process known as pion decay. The charged pion transforms into a muon ( $\mu$ ) of the same charge, along with the corresponding neutrino ( $\nu$ ) or antineutrino ( $\bar{\nu}$ ).


Hadronuclear interactions:



Two protons interact to form a neutral pion that cascades into two gamma ray photons.



Since a pion is made up of three particles: neutral, positively charged, and negatively charged, the pion produced decays in three stages. The first, as shown above, eventually releases a muon neutrino, positron, electron neutrino, and muon antineutrino.



The third stage results in a muon antineutrino, electron, electron antineutrino, and muon neutrino.

## Compton and Synchrotron Scattering:

Along with neutrino emissions, photons of various frequencies are emitted. There are two main processes that contribute to the energy of photons released by blazars:

First, Inverse Compton scattering occurs when a high-energy charged particle, usually an electron, interacts with a photon and transfers its energy to the photon (typically X-rays or gamma rays), resulting in a recoiling electron with diminished energy.<sup>9</sup> Second, synchrotron radiation which occurs as a charged particle is accelerated in a circle by magnetic fields. The charged particle, usually an electron again, releases a photon.<sup>10</sup> Detecting these photons in parallel with neutrino emissions gives scientists a way to correlate wavelengths such as X-rays, gamma rays, and radio waves with neutrino fluxes.

## Glashow Resonance:

Tying together the concepts of neutrinos and antineutrinos as well as neutrino production by astrophysical sources is a phenomenon called Glashow Resonance, a concept with more implications that are covered later in the paper.

Glashow resonance is a phenomenon predicted 60 years ago by Nobel laureate physicist Sheldon Glashow.<sup>11</sup> It explains that a high-energy electron antineutrino, with estimated energy above 6.3 PeV, can produce a W- boson when it interacts with an electron. No terrestrial accelerators can accelerate neutrinos to such high energies. This resonance is significant in the context of neutrino detection from BL Lac and FSRQ blazars due to the potential production of these high-energy antineutrinos in such environments. This means that only astrophysical neutrinos such as those created by blazars are capable of causing Glashow resonance. Secondary muons that cascade from the W- boson are what IceCube detects after Glashow resonance.

Glashow resonance provides validation of the standard model, a method of distinguishing neutrinos from antineutrinos, and a way to identify astronomical accelerators that produce neutrinos via hadronuclear or photohadronic interactions, with or without strong magnetic fields.

## IceCube - The World's Largest Cherenkov Telescope:

Detecting neutrinos is no simple feat. This is where the IceCube Detector comes in, the world's largest optical Cherenkov telescope. It is located in the Amundsen-Scott South Pole Station, made of one cubic kilometer of Antarctic ice. IceCube is designed to detect astrophysical and atmospheric neutrinos. Neutrinos have to interact with matter in order to be detected, a phenomenon that is very rare, given that neutrinos do not interact with the electromagnetic and strong force. To maximize the chance of a neutrino

interacting with matter, IceCube encompasses a huge area of 250 acres. A billion metric tons of clear ice with 5000 detectors allows huge amounts of light to be taken in.<sup>12</sup>

To ensure that sunlight cannot reach the sensors and prevent photon interference from any light that does not come from neutrinos, IceCube is also drilled 2.5 km deep. It is made up of 86 strings, 60 phototubes, and 5160 optical sensors.

When high energy particles interact with the ice, they create a shower of particles that travel faster than the speed of light in ice, though slower than the speed of light in a vacuum. Similar to a supersonic spacecraft emitting a sonic boom, these electrically charged particles release a cone of light known as Cherenkov radiation. This radiation is useful as it tells how much energy was deposited, which indicates how much energy the particle had, roughly where the particle came from, the type of particle, and possibly even the type of collision. In the ice, digital optical molecules (DOMs) designed as strings form much of the detector. With a photomultiplier and electronics, the DOMs detect Cherenkov light. Sensors collect the light, which is digitalized and time stamped.

The surface array of IceCube, called IceTop, is made up of a veto and calibration detector. IceTop detects neutrino air showers from cosmic rays in the energy range from 300 TeV to 1 EeV. Eight DOM strings packed more compactly within the ice are known as DeepCore. This component lowers the neutrino energy threshold to 10 GeV for closer atmospheric neutrinos. The IceCube lab processes data from the DOMs for information on what direction the neutrinos arrived from.

IceCube is not designed for low-energy neutrinos such as ones from the radioactive decay of rock, from sun, and from supernovae explosions. These all fall in the energy spectrum of thousandths GeV to a few hundredths GeV. Instead, IceCube commonly observes atmospheric neutrinos created by high energy particles that collide with nuclei and create a shower of hadrons that decay into neutrinos, as well as astrophysical neutrinos.

To contribute to multi-messenger astrophysics and validate the discoveries of IceCube, other telescopes are used to detect optical, radio wave, or gamma ray fluxes at the same time an astrophysical neutrino is detected. IceCube's most important collaborations include Laser Interferometer Gravitational-wave Observatory (LIGO), NASA's Fermi-gamma telescope, Major Atmospheric Gamma Imaging Cherenkov Telescope (MAGIC), Max Planck Institute for Radio astrophysics (MPIfR) in Germany, and Tracking Active Galactic Nuclei with Austral Milliarcsecond Interferometry (TANAMI).<sup>13</sup>

## IceCube Historical Data:

Since 2010, IceCube has made several groundbreaking discoveries in the realm of neutrino physics. Between 2010 and 2012, IceCube detected 26 high energy particles.<sup>14</sup> Their energies were more than one million times greater than those from supernovae in the Magellanic Cloud detected in 1987. These particles first indicated the possibility of neutrinos from outside the solar system.

Soon after, in April of 2012, a pair of neutrinos nicknamed Bert and Ernie were detected. Each one had an energy of above one PeV. This provided confirmation for neutrinos from outside the solar system and earned the Physics World 2013 Breakthrough of the Year.

In December of 2012, Big Bird was detected: a neutrino with less than a millionth the mass of an electron but energies a million million times that of an X-ray. With the energy of 2 quadrillion EeV, Big Bird was



the highest energy neutrino of its time and now still stands as the second most energetic. IceCube collaborated with scientists from the Fermi-Gamma ray telescope to connect Big Bird to the blazar PKS B1424-418. Observations showed that the blazar shone 15 to 30 times brighter than normal during the neutrino flare.

In July of 2018, IceCube for the first time confirmed the source of a high energy neutrino. By turning telescopes to observe the unusually bright bursts of energy from the blazar TXS 0506+056, scientists tied this blazar to the high energy neutrino event, named IceCube-170922A.

## Results and Discussion

In this section, we discuss more specific data from IceCube and other multi-messenger sources that demonstrate the importance of blazars in neutrino physics and implications of the data we receive from them.

### Proof that Neutrinos Originate from Blazars:

On September 22, 2017, IceCube detected a high-energy neutrino (estimated to be  $\sim 290$  tera-electron volts (TeV), and sent messages to other observatories around the globe. The Fermi Large Area Telescope Collaboration observed and identified a gamma ray source 0.1 degrees from the direction of the neutrino. The source, blazar TXS 0506+056, was observed to be in a flaring state. (The emission of neutrinos can be linked to gamma ray radiation, as explored later in this paper).<sup>15</sup>

The MAGIC Telescope Collaborations observed that the gamma-ray flux reached 400 GeV at times. TXS 0506+056 has a redshift of 0.34<sup>14</sup>. As depicted in Figure 2, the regions containing 50% and 90% of the neutrino event IceCube-170922A (indicated by dashed red and solid gray contours, respectively) are superimposed on an optical V-band sky image. Known gamma-ray sources within this area, previously identified by the Fermi spacecraft, are denoted by blue circles. The size of the circles corresponds to the 95% positional uncertainty of each source, and they are labeled with their respective names.




Figure 2: Circles represent positional uncertainty of previously detected gamma ray sources enabling multimessenger astrophysics to hone in on TXS 0506+056.<sup>14</sup>

The IceCube neutrino event aligns with the blazar TXS 0506+056, indicated by the pink square. The yellow circle represents the 95% positional uncertainty of very-high-energy  $\gamma$ -rays detected by the MAGIC telescopes during the subsequent observation campaign. Inset is a zoomed-in view of the region around TXS 0506+056, superimposed on an R-band optical image of the sky.

Scientists Gao et al., 2019 employed redshift as a basis to calculate the muon-neutrino luminosity (rate at which muon neutrinos are emitted from a source, measured in units of muon neutrinos per unit of time) and found that it matched the luminosity of the observed gamma-rays from that source.<sup>16</sup> These findings attribute the high-energy neutrino to the blazar TXS 0506+056 with a statistical significance of 3 sigma, or 99.7% confidence level.

Motivated by this discovery, the IceCube collaboration examined lower-energy neutrinos detected over the previous several years, finding an excess emission at the location of the blazar. In order to understand this, we will next explore the two subclasses of blazars: BL Lac Objects and Flat-Spectrum Radio Quasars (FSRQs).

### BL Lac Objects vs. Flat-Spectrum Radio Quasars:

FSRQs contain bright optical and UV emission lines, whereas BL Lac Objects lack these. FSRQs have a very high luminosity, but low source density (Figure 3).<sup>17</sup>




Figure 3: FSRQs are located in the upper left corner of this density-luminosity graph, illustrating that they are the highest in luminosity but lowest in source density of possible neutrino sources.<sup>17</sup>

Given the relatively small number of FSRQ sources in close proximity to each other, the detection of multiple neutrino samples from a single FSRQ source appears to be a reasonable expectation. However, this has not been detected, which indicates that FSRQs are likely not strong neutrino emitters.

This makes it unlikely for FSRQs to be strong neutrino emitters, because with such high power and low amount of FSRQs, one would expect to find multiple neutrino samples from a single FSRQ in the current data. However, since this is not the case, it seems unlikely that FSRQs would be strong neutrino emitters.



The FSRQ jet composition is also primarily made up of electrons and positrons (the antiparticles of electrons), whereas BL-Lacs are primarily electrons and protons, allowing for protons to be accelerated in BL Lacs. Photons are necessary for photohadronic interactions. Furthermore, neutrinos from astrophysical sources are expected to fall between the 0.1 -10 PeV, based on known relationships between the target photons, protons, and neutrinos.

However, the proton frequency of FSRQs corresponds to very high energies of protons, and therefore also high energies of neutrinos above the PeV threshold. This provides another possible reason for a lack of neutrino emission from FSRQs.

### Radiatively Inefficient Accretion (RIAF) flows and Blazars:

All super massive black holes have an accretion flow of particles around them. The quantity that determines the properties of this accretion flow is the accretion rate (the rate at which matter falls into the black hole).

A radiatively efficient flow has an accretion rate greater than  $10^{-2}$  Eddington units. This standard accretion disk is geometrically thin, optically thick (emits a wide range of electromagnetic radiation), and radiatively efficient. However, radiatively inefficient accretion flow is produced when the mass accretion rate is less than  $10^{-2}$ . In a radiatively inefficient accretion flow (RIAF), proton temperatures remain much hotter than in standard accretion disks. This high temperature causes the electrons to be stripped from the atoms, which creates a plasma. Plasma is very good at scattering photons, so the photons can travel through the RIAF without interacting with the matter, thus making RIAF geometrically thin and optically thick.<sup>18</sup>

The Spectral Energy Distribution (SED), or the range of energies emitted by the object at different wavelengths, of RIAFs is much broader than that of standard accretion flows.

The SED of FSRQ and BL Lacs is measured as how much energy is being emitted at different wavelengths due to both the jet and the accretion process. RIAFs are thought to be target photon sources for accelerated protons in these jets, resulting in the production of neutrinos (proton collisions with photons result in neutrinos), thus making BL Lac Objects to be strong candidates for neutrino emissions. Moreover, BL Lac objects are subdivided into three classes:

First, Low-Synchrotron-Peak BL Lacs (LBLs), such as TXS 0506+056, have a high density of RIAF photons in the soft X-ray range. This allows neutrino luminosity of LBLs to be 4 orders of magnitude larger than HBLs. In this class, the external RIAF photons are completely dominant. Second, in Intermediate-Synchrotron-Peak BL Lacs (IBLs), external RIAF photons are still more important than internal photons, though internal photons do have a presence. Third, in High-Synchrotron-Peak BL Lacs (HBLs), internal photons are the most significant, while external RIAF photons have little contribution. This means HBLs have a very low neutrino production efficiency. Figure 4 illustrates the contribution of internal and external photons for each subclass.<sup>18</sup>




Figure 4: Internal photons (solid lines) and external photons (dashed lines) contributing differently to their photon target fields in neutrino creating photohadronic processes.<sup>18</sup>

BL Lac objects, especially LBLs, with weak emission lines and a lack of external radiation fields, were originally thought to be less likely as high-energy neutrino candidates. However, the high-energy neutrino from TXS 0506+056 came from a LBL. Despite HBLs emitting stronger X-rays, high-energy neutrinos have not been detected from them.

With evidence of RIAFs allowing for neutrino production and the previously mentioned characteristics that prevent FSRQs from being strong neutrino emitters, it can be concluded the BL Lacs are much more likely to be a major source of astrophysical neutrinos.

### Spectral Energy Distribution (SED) of Blazars:

The Spectral Energy Distribution (SED) of all blazars come in two categories. The low-energy category, from radio to UV are likely due to synchrotron radiation by relativistic electrons (electrons that approach the speed of light). The high-energy category, from X-rays to gamma rays, is a result of the same relativistic electrons undergoing inverse Compton scattering off target photons or other hadronic processes. Dividing the broad band SED into two categories, the low-energy category and the high-energy category, is called the leptonic model for modeling electromagnetic radiation from blazars.<sup>18</sup>

Blazars are further subdivided into three classes depending on the highest point of synchrotron emission: low-synchrotron peaked, intermediate-synchrotron peaked, and high-synchrotron peaked. While FSRQs only fall under the low-synchrotron peaked class, BL Lac objects can fall under any of the three classes.

As we have previously seen in this paper, the production of neutrinos requires hadronic processes, thus causing the leptonic model to be modified. The hadronic model for electromagnetic radiation describes how cosmic-ray nucleons at energies  $\sim 10$  PeV interact with UV photons to produce charged and neutral pions. Neutral pions decay into gamma-ray photons while charged pions decay into a muon and electron as well as neutrinos that travel to Earth.

However, this hadronic model does not entirely match scientific observation. X-ray emissions by the secondary electrons predicted in the model overshoot the actual observed X-ray flux. This means that the ideal model for electromagnetic radiation from blazars should not be entirely the hadronic model. A hybrid model was proposed to allow for neutrino emission but constrain the X-ray flux.<sup>16</sup> This hybrid

model says that most photons are of leptonic origin, and any contribution by the hadronic processes is limited so that there are not so many X-rays produced that they overshoot the flux constraints. The hybrid model is shown in Figure 5.




Figure 5: Hybrid model of neutrino production incorporating leptonic and hadronic parts, without overshooting the observed X-ray flux.<sup>16</sup>

Another constraint put on this model includes the Eddington limit, meaning that radiation from the blazar cannot be so powerful that it counteracts the force of gravity and blows away surrounding matter. The maximum contribution from photohadronic interactions is constrained by the fact that efficient neutrino production requires high energy photons, electrons, and positrons. This means that the electromagnetic cascade has to be visible as X-rays and TeV gamma rays.

As shown in Figure 6, Time-dependent modeling of emissions before, during, and after the injection boost of a blazar indicates a sharp increase in amplitude of hard X-rays and gamma-rays during the neutrino flux.




Figure 6: Relationship between hard X-rays, gamma-rays, and neutrinos underscoring the importance of multiwavelength astrophysics.<sup>19</sup>

## Multiwavelength Astrophysics:

In multimessenger or multiwavelength astrophysics, studying all the processes that contribute to the overall observed electromagnetic spectrum is significant. Information from IceCube led to a spur in multi-messenger astrophysics as other detectors sought to correlate neutrino emission with other types of radiation. By studying the emissions across different wavelengths such as gamma rays, X-rays, radio waves, etc, scientists can piece together a more comprehensive understanding of the astrophysical phenomena and the underlying physical processes at play.

### Radio Waves:

The Owens Valley Radio Observatory (OVRO) further analyzed radio data from blazars, detecting a long-term outburst of radio waves before reaching a peak during neutrino emission for both PKS 1502+106 and TXS 0506+056. Based on this and other studies, the correlation between IceCube neutrinos and radio-bright blazars has been quantified at a p-value of 0.002, indicating that the correlation between IceCube neutrinos and radio-bright blazars has a 0.2% probability of it being due to pure chance.<sup>19</sup>

Despite this statistical significance, observations from OVRO still did not indicate a strong correlation between radio emission and neutrinos since this was only two out of the five sources observed by OVRO. A third source was observed only after 70 days of the arrival of the neutrino. Though all five sources are monitored in radio, none demonstrated long term radio wave activity, indicating that more observation is needed to correlate them to the emission of high energy neutrinos.

On the other hand, a more recent study found that in recent years, an increasing number of neutrinos detected can be correlated to radio blazars.<sup>20</sup> It was found that radio blazars significantly correlated with the entire 7-year dataset from IceCube, including radio waves from lower energies as low as a few TeV range. Combined with other analyses on independent data, the correlation of neutrinos with radio waves below and above 200 TeV was found to be significant at 4.3 standard deviations, which indicates that the differences between groups of data is large enough for there to be statistical significance. This means that there was enough variability in the evidence that the correlation was strong enough to overcome external factors.

With contrasting data from multiple analyses, more data needs to be collected to confirm a correlation between radio emission and neutrino emission. Recent data may be more accurate as it uses Very Long Baseline Interferometry (VLBI), a process that compares the distances traveled by light to each telescope in a system of telescopes placed far from each other. This allows for highly accurate astronomical data, especially when focusing on radio blazars to relate them to neutrino emission. The evidence indicates there is a high likelihood of neutrinos originating from radio blazars, but more monitoring by observatories such as OVRO needs to be done to identify the chance of a flux of radio waves during a single high-energy neutrino event.

### Gamma rays:

Gamma rays were also correlated with neutrino emission. The first likely extragalactic neutrino counterpart is the gamma-ray blazar TXS 0506+056, identified on September 22, 2017. The gamma-ray blazar was found to be in a flaring state in spatial and temporal coincidence with the arrival of the 290 TeV neutrino event IC-190722A.<sup>19</sup>

Since TeV and PeV photons are quickly absorbed through photon-photon annihilation and cascade down to lower energies, GeV gamma-rays are the closest in energy to high-energy neutrinos. This makes them important to neutrino detection.

In the study of gamma-rays from astrophysical bodies, there were two neutrino observations from the same source.<sup>21</sup> The source emitted a 160-day-long neutrino flare in 2014 and again in 2015. It was not found to be accompanied by an increased activity in gamma-ray, optical, or radio wavelength. It was discovered that the gamma-ray spectrum hardened, but the discovery was not statistically significant. One study found that the hardening of the gamma-ray spectrum was statistically compatible at less than 2 standard deviations with the average spectrum, meaning there was likely no significant increase of the gamma-rays due to neutrino events.

Follow-up studies to discover other neutrino sources were initiated. However, while only measuring neutrino data and ignoring multiwavelength data, there was no detection of flares above the expected atmospheric background. This suggests that neutrino emission from these sources may be accompanied by emission at other wavelengths, such as radio waves, gamma rays, or X-rays. It is also possible that the neutrino emission from these sources is very weak and difficult to detect.

The Fermi-LAT telescope is sensitive to gamma rays, making it a crucial collaborator with Ice Cube in the detection of astrophysical neutrinos. The study of more blazars, this time focusing on multi-wavelength data, reveals more information about gamma-rays. Below is the analysis of data from two blazar neutrino sources.<sup>21</sup>

The study of the blazar 1H 0323+342 reveals that this source has a p-value of 0.08 for gamma-ray fluxes during neutrino flares, whereas other sources show p-values from 0.17 to 0.92. The neutrino from 1H 0323+342 came during a small increase in the gamma-ray flux, while an increase in UV, X-ray, and optical emission came one month earlier.

The blazar PKS 1502+106, a possible neutrino source for the neutrino event IC-190730A, is particularly interesting to scientists because it is extremely gamma-ray bright: it is the 15th brightest out of 2863 in terms of gamma-ray luminosity. The gamma-ray emission from this blazar has been explained to be from interactions between jet electrons and photons from the broad line region. This information indicates that the BLR photons could provide a target photon field for hadrons in the jet, allowing for neutrino production.

Over an 11 year period, the gamma-ray emission from PKS 1502+106 has been studied, identifying periods of both quiet and flaring states. Altogether, two hypotheses are consistent with the observed data. The first is that the gamma-ray brightness of blazars is linearly correlated with neutrino emission. This is supported by the neutrino event IC-170922A being correlated with a bright gamma-ray flare from TXS 0506+056.

However, blazar stacking analyses show that if the neutrino spectrum follows an  $e^{-2.5}$  power law, the gamma-ray blazar contribution is limited to less than 27%. If the neutrino follows a steeper power law of  $e^{-2}$ , the stacking analysis shows that contribution from the gamma-ray blazars expands to become 40-80%.<sup>19</sup> Figure 7 illustrates the difference between these power laws.




Figure 7: The graph showing the two possible power laws that the neutrino spectrum can follow:  $e^{-2.5}$  and  $e^{-2}$ . (Author).

The power law determines the relationship between the flux of neutrinos at different energies as the output of the energy of the neutrino. Stacking analyses show that if the neutrino flux follows a  $e^{-2}$  power law, the contribution by gamma-ray blazars expands to 40-80%. Note that this applies to neutrinos between the 10 TeV to 100 TeV range, while this graph is only used to depict the general pattern of these two power laws.

The second hypothesis, the background hypothesis, is that neutrino emission is not correlated to gamma-ray emission. There are a few possible reasons for this. The gamma-ray flux that has been identified a few times with neutrino emission could be coincidental. Another reason that scientists might conclude that neutrino emission does not have a correlation with gamma-rays is that gamma-rays from some neutrino sources are produced in optically thick regions, so gamma-rays could be absorbed and then cascade down to X-rays.

If the first hypothesis is true, gamma-ray flare coincidences can be used to study neutrinos and cosmic-ray acceleration. This also allows us to associate high-energy neutrino sources with more distinct properties. Since it would be impossible to detect the production of gamma rays beyond the optically thick regions, more observations would be needed to collect data points of various blazars with gamma-ray emission. IceCube-Gen2 will be especially useful for this as it will also be much more sensitive to weaker neutrino sources.<sup>22</sup>

## X-rays:

The association of X-rays with neutrino producers is the most statistically significant compared to the other correlations discussed thus far. As explained in the previous section, broadband SED modeling suggests that X-ray photons from the corona (the outer edges of the blazar) could make up a target photon field for photohadronic interactions that produce high-energy neutrinos. Figure 8 shows a



time-dependent model of the different parts of a Spectral Energy Distribution shows an increase in the flux of Hard X-rays and TeV gamma-rays at the same time there is a flux in muon neutrino events.<sup>19</sup>




Figure 8: Time-dependent model of the Spectral Energy Distribution (SED) revealing concurrent increase in the flux of Hard X-rays and TeV gamma-rays coinciding with a flux in muon neutrino events.<sup>19</sup>

The steepening of the graph of the gamma ray spectrum is also explained by X-ray coronal photons because they absorb gamma-rays through pair production.

An analysis of X-ray neutrino sources was conducted by cross-matching X-ray sources with a sample set of Very-Long-Baseline-Interferometry (VLBI) bright sources, which are primarily blazars. If telescopes detected X-ray emission within a certain threshold of a VLBI source, it was named an X-ray source.<sup>20</sup> Then, an analysis was completed that counted the matches between neutrinos and X-ray VLBI blazars. This was compared to a count of the random expectations of the matches. For one sample, SRG/ART-XC catalog, the statistical significance of neutrinos correlated to X-ray VLBI blazars was 0.5%. In fact, all three of the hard X-ray catalogs studied, SRG/ART-XC, Swift/BAT, and INTEGRAL/IBIS reveal a correlation with IceCube neutrinos.

However, there was no correlation within the ROSAT catalog, a set of soft X-ray blazars. Thus, it was found that blazars that emit neutrinos likely have the characteristic of X-rays above 10 keV.

X-rays are found to make up the target-photon field for photohadronic interactions responsible for neutrino emission, given that they are produced in the jet. Thus, the emission of hard X-rays also explains the correlation of radio waves with neutrino emission because the number of synchrotron-self-Compton photons is proportional to the amount of radio-waves emitted by the photons. These synchrotron-self-Compton photons are located in the jetstream of a blazar.

Additionally, higher energies of GeV gamma ray photons are found to interact with low-energy photons and produce electron-positron pairs and cascading down to lower energies, which lose efficiency in the

GeV range. So, although gamma-ray photons may accompany neutrinos in production, observing X-rays is a more reliable method to trace neutrino production.

X-ray flares are important for neutrino emission because they can produce pions, which then decay into neutrinos. Pions are produced when high-energy protons interact with photons, and X-ray photons are energetic enough to produce pions. Additionally, X-ray fluxes are often optically thick, which means that there is a high probability of protons interacting with photons. One model explains why focusing on X-ray flares is important. Since X-ray photons are energetic targets for photomeson interactions (interactions between high-energy protons and photons), they reduce the energy required of protons for pion production. Pions are unstable particles that decay into other particles, including neutrinos.<sup>23</sup> Thus, X-rays have the highest statistical significance in being correlated with neutrino emission not just based on observational analysis but also because they explain crucial production processes for neutrinos.

Therefore, as analyzed in this paper thus far, multiwavelength data is crucial to the understanding of neutrino production from blazars.

## Neutrinos, Antineutrinos, and Glashow Resonance:

Much is still unknown about astrophysical neutrinos. Certain neutrino events have been traced back to their sources, allowing for there to be more data on blazars and neutrino production processes. However, other neutrino events have not been associated with a specific astrophysical source, but still provide crucial information. In 2016, IceCube measured a shower of particles with energies  $6.05 \pm 0.72$  PeVas, determined by Cherenkov radiation. Antineutrinos Interacting with electrons in the ice results in the formation of a W boson, which decays into secondary muons. The detection of these secondary muons by IceCube indicated the occurrence of Glashow Resonance.<sup>11</sup>

The huge number of electrons located in Earth allows for an increased chance of Glashow resonance occurring. This is significant as it allows antineutrinos to be distinguished within the neutrino flux. Now, Glashow resonance can occur between either a neutrino and a positron (antielectron) or an antineutrino and electron, but because the Earth is made of matter and not antimatter, the neutrinos will almost never hit a positron.<sup>23</sup> However, as there is an abundance of electrons in everyday environments, some antineutrinos will hit electrons. So, if evidence of Glashow resonance is detected, scientists can be reasonably certain that this is evidence of an antineutrino.

Now, in a nucleon (the collective term for protons and neutrons), valence quarks are the primary quarks that contribute to its composition as well as being involved in the strong force interactions that hold nucleons together (akin to valence bonds in atoms). For example, a proton is made up of two up-quarks and one down-quark. A neutron is made of two down and one up-quark.<sup>25</sup>

Valence quarks are distinct because they are the only particles within a nucleon that are unpaired with an antiquark. Antiquarks are the antimatter counterparts of quarks. Also, valence quarks are left-handed (their spin is aligned to their momentum) the weak force acts on left-handed particles and right-handed anti-particles. In particle physics, quarks come in left- and right-handed versions. Only the left-handed ones can interact with W bosons because they have weak isospin, similar to electric charge for electromagnetic interactions. Due to this, valence quarks are more likely to engage with neutrinos and antineutrinos in comparison to other types of quarks within the nucleon. In other words, in astrophysical contexts, nucleons serve as interaction targets for neutrinos and antineutrinos.

However, at higher energies, the ratio of other quarks to valence quarks becomes high enough that valence quarks become lost in the noise. This is why in the realm of high-energy neutrino interactions,

Glashow resonance, which is primarily driven by interactions with valence quarks, becomes increasingly important and thus proving high-energy electron antineutrino collisions.

Prior to the new analysis of neutrinos and antineutrinos, studies of the astrophysical neutrino flux assumed that there was an equal contribution of neutrinos and antineutrinos, and they focused only on a three-flavor composition. The three-flavor composition assumes that there are three types of neutrinos: electron neutrinos, muon neutrinos, and tau neutrinos. The ratio of these three flavors is given by  $(f_e : f_\mu : f_\tau) = (1:2:0)$ , meaning that there is one electron neutrino for every two muon neutrinos and no tau neutrinos.<sup>24</sup>

However, it was shown that this assumption is not correct. He showed that neutrino production by high-energy sources results in an asymmetry between the particles, necessitating a six-flavor composition. The six-flavor composition includes the three neutrino flavors and their corresponding antineutrino flavors.

This asymmetry is caused by the different ways that different hadronuclear and photo-hadronic interactions produce positively charged pions and negatively charged pions.<sup>24</sup> Positively charged pions decay to produce electron neutrinos and muon neutrinos, while negatively charged pions decay to produce electron antineutrinos and muon antineutrinos.

As a result of this asymmetry, the ratio of neutrinos to antineutrinos in the astrophysical neutrino flux is not equal. New ratios for the photo-hadronic scenario and the hadronuclear scenario were calculated. This ratio is given in the form of  $(\{f_e, -v_e\} : \{f_\mu, -v_\mu\} : \{f_\tau, -v_\tau\})$ , with  $-v_e$  representing an electron antineutrino,  $-v_\mu$  representing a muon antineutrino, and  $-v_\tau$  representing a tau antineutrino. The photohadronic ratio is  $(\{1, 0\} : \{1, 1\} : \{0, 0\})$ , and the hadronuclear ratio is  $(\{1, 1\} : \{2, 2\} : \{0, 0\})$ .

As explained before, Glashow resonance is a unique interaction between a high-energy electron antineutrino and an electron. It results in the creation of a  $W^-$  boson. This resonance process is significant because it enables the differentiation between neutrinos and antineutrinos in the astrophysical flux. Liu's work highlights the necessity of a six-flavor model to account for the intrinsic asymmetry between neutrinos and antineutrinos produced by high-energy sources interacting with earthly electrons. This six-flavor model encompasses both neutrino and antineutrino counterparts of the original three flavors (electron neutrinos, muon neutrinos, and tau neutrinos). This is significant as the processes by which neutrinos were produced, photohadronic or hadronuclear, can be distinguished based on the ratios of the neutrinos and antineutrinos detected in a flux.

In essence, this research unveils the underlying mechanisms that lead to an imbalance between neutrinos and antineutrinos, a possible evidence of CP violation. Glashow resonance complements this understanding by providing a means to identify electron antineutrinos, which are vital components of this newly recognized asymmetry. The resonance process, occurring due to the presence of electrons, allows for the discrimination between neutrinos and antineutrinos, thus enriching our insights into astrophysical neutrino production processes. This is the beginning of a deeper exploration of how neutrinos can provide evidence of CP violation.<sup>26-27</sup>

## Conclusion

This review paper critically examined various hypotheses regarding neutrino physics and provides insights into the most promising ones. It also highlighted the limitations of existing experiments and

underscores the inadequacy of the standard model in addressing fundamental questions about the universe, particularly concerning neutrinos.

IceCube has been the first neutrino detector to relate a neutrino to an astrophysical source, opening the realm of multi-messenger astrophysics.

IceCube has answered key questions about high energy neutrinos. It correlates neutrinos to blazars as sources with a statistical significance of  $3.5 \sigma$ . The Radiatively Inefficient Accretion Flow (RIAF) that likely exists in the nucleus of a BL Lac Object provides a strong region for neutrino production, compared to a standard accretion flow of FSRQs.

Neutrinos provide crucial information about astrophysical bodies. Because neutrinos can be associated with blazars, we can provide specific characteristics of blazars. Photohadronic and hadronuclear interactions, crucial in neutrino production, require accelerated protons and a photon target field. As seen in the paper, blazars must provide these components in the form of jet streams, RIAFs, and possibly X-ray target photon fields from the jet stream.

To identify neutrino emission from astrophysical objects, IceCube data reveals that hard X-rays are key. The correlation between neutrinos and gamma-rays or radio-waves is yet to be statistically significant and more data needs to be collected. X-rays provide an ideal target photon field for neutrino production.

IceCube is beginning to answer questions about how CP violation can explain the matter-antimatter imbalance in the universe. Glashow resonance provides a way to distinguish between neutrinos and antineutrinos originating from astrophysical sources, but more data needs to be collected to see if this explains CP violation.

Improvements on IceCube are planned to be completed by 2032 to create IceCube-Gen2. With the installation of 750 advanced photodetectors and calibration devices, IceCube-Gen 2 will increase the volume of IceCube by tenfold, expanding it to 10 cubic kilometers. Its higher sensitivity and sharper focus will allow it to measure super PeV events and above 100 TeV astrophysical flux. It will be the first to measure neutrino cross sections at PeV and EeV energies, well beyond the capabilities of particle accelerators.

When it comes to multi-messenger astrophysics, data collected by IceCube-Gen2 will help provide answers to some of the uncertainties presented in this paper, such as the correlation of neutrino emission with radio and gamma-ray fluxes. IceCube-Gen2 will be able to cover a range of energies, with optical and radio detectors providing continuous coverage of emissions across the spectrum. This will provide more data that can then be used to correlate radio waves and gamma rays to neutrino events. IceCube-Gen2 will also be much more sensitive to weaker neutrino sources, up to five times fainter, than what could be detected with IceCube. As a result, it will be sensitive to hidden sources in optically thick environments where only neutrinos escape.

Being able to analyze this multimessenger information from high-energy astrophysical neutrinos is a vital tool that IceCube provides in understanding key processes in the universe.

## Acknowledgements

I would like to thank my mentor, Jaclyn Schillinger, for her extensive knowledge and helping me understand the technical aspects of neutrino physics and expand my knowledge.

## References

1. Sundermier, A. Symmetry magazine. Game changing neutrino experiments page. <https://www.symmetrymagazine.org/article/game-changing-neutrino-experiments> (published 2018-05-24).
2. Harris, D. US Department of Energy Office of Science (Fermilab), DOE explains Neutrinos page. <https://www.energy.gov/science/doe-explainsneutrinos>. (published 2020-06)
3. Aartsen, M.G.; Abbasi, R.; et al (The IceCube Collaboration). Detection of a particle shower at the Glashow resonance with IceCube. *Nature*. 2021, 591, 220-224 (2021). DOI: 10.1038/s41586-021-03256-1
4. Piergrossi, J. Symmetry magazine. Neutrinos, the Standard Model misfits. <https://www.symmetrymagazine.org/article/february-2013/neutrinos-the-standard-model-misfits/> (published 2013-02-13)
5. Reissman, H. Ideas.TED.com page, Could the neutrino hold the answers to some of the universe's big questions. <https://ideas.ted.com/could-the-neutrino-hold-the-answers-to-some-of-the-universes-big-questions/> (published 2017-05-05)
6. Fermi National Accelerator Laboratory, Cosmic neutrinos page <https://neutrinos.fnal.gov/sources/cosmic-neutrinos> (published 2020-05)
7. Baird, C West Texas A&M Science Questions with Surprising Answers page. How does a black hole give off light? <https://www.wtamu.edu/~cbaird/sq/2016/03/29/how-does-a-black-hole-give-off-light/> (published 2016-03-29)
8. Liu, R.Y. ; Wang, K. ; Xue, R. ; Taylor, A.M.; Wang, X.Y.; Li, Z. ; Yan, H. Hadronuclear interpretation of a high-energy neutrino event coincident with a blazar flare. 99, *Physical Review D* 99, 063008 (2019), DOI: 10.1103/PhysRevD.99.063008.
9. Mushotzky, R. University of Maryland, Dept. Of Astronomy page. [https://www.astro.umd.edu/~richard/ASTR480/Beckmann\\_Longair\\_Radiation3.pdf](https://www.astro.umd.edu/~richard/ASTR480/Beckmann_Longair_Radiation3.pdf) (published 2019-08-28, accessed 2023-07-09)
10. NIST.gov Sensor Science Division page. What is synchrotron radiation? <https://www.nist.gov/pml/sensor-science/what-synchrotron-radiation>. (published 03-02-2010, Updated 2021-06-02)
11. Aartsen, M.G.; Abbasi, R. et al (The IceCube Collaboration). Detection of a particle shower at the Glashow resonance with IceCube. *Nature*. 2021, 591, 220-224 (2021). DOI: 10.1038/s41586-021-03256-1
12. University of Wisconsin-Madison, ICECUBE Neutrino Observatory page. <https://icecube.wisc.edu/science/icecube/> (accessed 2023-07-13).





13. Strassler, M. IceCube: A Giant Frozen Neutrino Catcher page. <https://profmattstrassler.com/articles-and-posts/particle-physics-basics/neutrinos/how-one-detects-neutrinos/icecube-a-giant-frozen-neutrino-catcher/>. (published 2013)
14. Tillman, N.T.; Space.com magazine page. IceCube: Unlocking the Secrets of Cosmic Rays. <https://www.space.com/41170-icecube-neutrino-observatory.html>. (published 2018-07-14)
15. The IceCube Collaboration, Fermi-LAT, MAGIC, et al. Multimessenger observations of a flaring blazar coincident with high-energy neutrino IceCube-170922A. *Science* 07-13-2018, Vol 361, Issue 6398 DOI: 10.1126/science.aat1378
16. Gao, S.; Fedynitch, A.; Winter, W.; Pohl, M. Modelling the coincident observation of a high-energy neutrino and a bright blazar flare. *Nature*. 2019-5-10. arXiv:1807.04275v3 DOI: 10.1038/s41550-018-0610-1
17. Kowalski, M. Status of High-Energy Neutrino Astronomy. *Journal of Physics: Conference. Series.* 632 012039 2014-10. DOI: 10.1088/1742-6596/632/1/012039
18. Righi, C.; Tavecchio, F.; Inoue, S. Neutrino emission from BL Lac objects: the role of radiatively inefficient accretion flows. *Monthly Notices of the Royal Astronomical Society: Letters*, Volume 483, Issue 1. 2019-02. Pages L127–L131. DOI: 10.1093/mnrasl/sly231
19. Franckowiak, A; Garrappa, S. et al. Patterns in the Multiwavelength Behavior of Candidate Neutrino Blazars. *The Astrophysical Journal*, The American Astronomical Society, 2020-04-28. DOI 10.3847/1538-4357/ab8307
20. Plavin, A.V.; Burenin, R.A.; Kovalev, Y.Y.; Lutovinov, A.A.; Starobinsky, A.A.; Troitsky, S.V.; Zakharov, E.I. Hard X-ray emission from blazars associated with high-energy neutrinos. *ArXiv INR-TH-2023-007*, 2023 September, DOI: 10.48550/arXiv.2306.00960
21. Garrappa, S. et al. Investigation of Two Fermi-LAT Gamma-Ray Blazars Coincident with High-energy Neutrinos Detected by IceCube. *The Astrophysical Journal*. Volume 880, Number 2. 2019-07-31. DOI 10.3847/1538-4357/ab2ada
22. The IceCube-Gen2 Collaboration, IceCube-Gen2: The Window to the Extreme Universe. *Journal of Physics*, G48 (2021) 6, 0605012020-08-10. 2021-04-29. arXiv:2008.04323v1 [astro-ph.HE]. DOI: 10.1088/1361-6471/abbd48
23. Stathopoulos, S.I.; Petropoulou, M.; Giommi, P.; Vasilopoulos, G.; Padovani, P.; Mastichiadis, A. High-energy neutrinos from X-rays flares of blazars frequently observed by the Swift X-ray Telescope. *Monthly Notices of the Royal Astronomical Society*, Volume 510, Issue 3, Pages 4063–4079, March 2022 DOI: 10.1093/mnras/stab3404.
24. Liu, L.; Ningqiang, S., and Vincent, A.C. Probing neutrino production in high-energy astrophysical neutrino sources with the Glashow resonance. *American Physical Society Journal. Phys. Rev. D* 108, 043022. 2023-07-24. DOI: 10.1103/PhysRevD.108.043022
25. Cates, G.; Wojtsekhowski, B.; Carter, K. US DoE Jefferson Lab Page. Quarks Pair Up in Protons (and Neutrons). <https://www.jlab.org/quarks-pair-protons-and-neutrons>. (published: n.a., accessed 2023-10-20)
26. University of Alabama, Department of Physics & Astronomy page. Antimatter Neutrino from Space Distinguished from Normal Matter Neutrino for First Time Using Unique Process. <https://physics.ua.edu/2021/03/10/antimatter-neutrino-from-space-distinguished-from-normal-matter-neutrino-for-first-time-using-unique-process/> (published: 2021-03-10)





27. Chatterjee, A.; Devi, M.M.; Ghosh, M; Moharana, R.; Raut, S.K. Probing CP violation with the three years ultra-high energy neutrinos from IceCube. Rev. D 90, 073003 (2014). American Physical Society, Phys. Rev. D 90, 073003. DOI: 10.1103/PhysRevD.90.073003

## Author

Avni Iyer is a sophomore at Leland High School, San Jose, CA, USA, with calculus, physics and engineering being her primary passions. She plans to specialize in aerospace engineering and applied astrophysics.